


Bangsamoro Autonomous Region in Muslim Mindanao



Water Availability for Corn
Prevailing Stage : None

Harvesting and/or
pre-planting
stage is possibly
ongoing.

Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Basilan	118.7	I	30.6	I	73.7
		II	27.6	II	50.9
		III	26.1	III	38.7
		IV	26.1	IV	50.2
Lanao del Sur	260.8	I	153.0	I	266.0
		II	142.3	II	205.9
		III	137.4	III	168.2
		IV	137.4	IV	204.0
Maguindanao	285.6	I	131.6	I	262.6
		II	122.2	II	191.1
		III	117.9	III	153.7
		IV	117.9	IV	189.1
Sulu	99.8	I	33.8	I	52.6
		II	30.5	II	35.7
		III	28.9	III	27.4
		IV	28.9	IV	35.3
Tawi-Tawi	167.6	I	76.4	I	154.0
		II	68.2	II	109.3
		III	64.6	III	83.5
		IV	64.6	IV	108.1

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of April

Rice CS:
(I) Nursery
(II) Vegetative
(III) Reproductive
(IV) Flowering


Corn CS:
(I) Establishment
(II) Vegetative
(III) Silking
(IV) Reproductive

CCI Category:
Inadequate
Sufficient
Excess


Regional Summary

Except for Basilan and Sulu, sufficient rainfall received for the month was favorable for the rice crops at any stage for the region. Inadequate rainfall for the month could have caused harm on rice crops at any stage in Basilan and Sulu. Meanwhile, the condition of crops in the farm areas were generally fair, according to the NDVI. The SPEI3 suggests that the region generally experienced near normal condition of accumulated rainfall in the past three months. The extreme rainfall indices show maximum 1-day (50-150mm) and 5-day (100-250mm) rainfall which may have caused harm on crops in areas of Lanao del Sur and Maguindanao. This extreme rainfall was primarily related to the Easterlies and Trough of LPA during the early week of the month.


NDVI



SPEI3 (Feb-Mar-Apr)



RX1day



RX5day

