



Region VIII (Eastern Visayas)



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice CS	Rice CCI (%)		Corn CS	Corn CCI (%)	
			CS	CCI (%)		CS	CCI (%)
Biliran	92.3	I	36.0	I	92.6	I	92.6
		II	31.4	II	58.1	II	58.1
		III	29.4	III	40.7	III	40.7
		IV	29.4	IV	57.1	IV	57.1
Eastern Samar	157.6	I	79.2	I	153.3	I	153.3
		II	71.7	II	116.0	II	116.0
		III	68.4	III	91.1	III	91.1
		IV	68.4	IV	114.9	IV	114.9
Leyte	96.7	I	35.1	I	88.9	I	88.9
		II	30.7	II	55.6	II	55.6
		III	28.7	III	38.4	III	38.4
		IV	28.7	IV	54.7	IV	54.7
Northern Samar	191.2	I	105.3	I	197.6	I	197.6
		II	96.2	II	141.2	II	141.2
		III	92.1	III	109.9	III	109.9
		IV	92.1	IV	139.6	IV	139.6
Samar	189.4	I	96.3	I	151.7	I	151.7
		II	88.2	II	112.6	II	112.6
		III	84.4	III	87.4	III	87.4
		IV	84.4	IV	111.4	IV	111.4
Southern Leyte	86.7	I	33.2	I	54.5	I	54.5
		II	30.0	II	35.2	II	35.2
		III	28.6	III	25.8	III	25.8
		IV	28.6	IV	34.7	IV	34.7

Crop Stage (CS) highlighted in black is the dominant stage during the month of April

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Flowering
Corn CS: (I) Establishment, (II) Vegetative, (III) Silking, (IV) Reproductive
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

Regional Summary

Except for the provinces of Biliran, Leyte, and Southern Leyte, which received insufficient amounts of rainfall during the month and may not be enough for the preparation of rice paddy. The corn crops in this Region, which were still in a young stage, received enough rainfall during the month.

The NDVI shows healthy vegetation over the region. SPEI3, showing the past three months rainfall, indicates that the observed rainfall was near normal to slightly wetter condition.

The RX1day and RX5day with maximum accumulated rainfall amounting to 100mm to 150mm, respectively are beneficial for the preliminary land preparation for rice paddy and also for standing young corn crops.

