



# Region XIII (Caraga)



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice CS	CCI (%)	Corn CS	CCI (%)
Agusan del Norte	106.3	I	40.1	I	92.7
		II	35.2	II	61.1
		III	33.1	III	43.1
		IV	33.1	IV	60.2
Agusan del Sur	105.0	I	36.8	I	81.7
		II	33.0	II	53.4
		III	31.3	III	39.0
		IV	31.3	IV	52.6
Dinagat Islands	133.9	I	53.5	I	84.4
		II	48.5	II	59.6
		III	46.2	III	44.5
		IV	46.2	IV	58.8
Surigao del Norte	127.8	I	52.9	I	98.9
		II	48.2	II	69.9
		III	46.1	III	54.2
		IV	46.1	IV	69.0
Surigao del Sur	126.4	I	59.3	I	123.9
		II	53.3	II	86.5
		III	50.6	III	65.1
		IV	50.6	IV	85.4

Crop Stage (CS) highlighted in black is the dominant stage during the month of April

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Flowering  
Corn CS: (I) Establishment, (II) Vegetative, (III) Silking, (IV) Reproductive  
CCI Category: Inadequate (yellow), Sufficient (green), Excess (blue)

## Regional Summary

Rainfall received for the month was adequate to support the corn crops at any stage in Surigao del Norte and Surigao del Sur. This sufficiency in rainfall was also beneficial for the corn crops at non-prevailing stage in Agusan del Norte, Agusan del Sur, and Dinagat Islands, but the corn crops at the prevailing Silking stage might have been damaged due to insufficient rainfall received for the month in these areas.

The crops in the farm areas were in good condition based on the NDVI. In addition, the SPEI3 suggests that the rainfall for the past three months was in near the normal condition throughout the region. The extreme rainfall indices do not show any significant heavy rainfall that might have brought damage on crops in Caraga region.

