

Region XIII (Caraga)

Water Availability for Rice
Prevailing Stage : None

Provincial Values


	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Agusan del Norte	123.2	I	48.7	I	112.6
		II	43.7	II	71.4
		III	41.5	III	52.6
		IV	75.8	IV	70.4
Agusan del Sur	150.3	I	92.5	I	196.8
		II	83.7	II	140.6
		III	79.7	III	108.3
		IV	133.7	IV	139.0
Dinagat Islands	96.4	I	49.0	I	87.0
		II	43.9	II	61.5
		III	41.6	III	44.0
		IV	71.3	IV	60.7
Surigao del Norte	125.2	I	67.4	I	128.2
		II	61.3	II	95.2
		III	58.5	III	75.0
		IV	93.2	IV	94.2
Surigao del Sur	181.3	I	98.5	I	208.0
		II	89.5	II	150.0
		III	85.5	III	114.3
		IV	141.3	IV	148.2

Crop Stage (CS) highlighted in bold is the dominant stage during the month of April

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity
CCI Category: Inadequate Sufficient Excess

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : (III) Reproductive



Regional Summary

Most parts of CARAGA Region received sufficient rainfall, except for the province of Agusan del Norte and Dinagat Islands which received inadequate rainfall for rice crops during pre-planting or harvesting activities. Majority of corn crops in the prevailing Reproductive stage in the region also received sufficient rainfall, except at the Dinagat Islands.

The SPEI3 further indicates that the Region experienced slightly wetter conditions over the past three months. While the RX1day shows no extreme 1-day rainfall, RX5day indicates significant accumulation of about 50-150 mm due to the effects of the Intertropical Convergence Zone (ITCZ), easterlies, localized thunderstorms, and Low-Pressure Area (LPA).

NDVI

NDVI satellite data not yet available.

SPEI3 (Feb-Mar-Apr)



RX1day



RX5day

