


# Bangsamoro Autonomous Region in Muslim Mindanao

Water Availability for Rice  
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn  
Prevailing Stage : (IV) Maturity



## Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Basilan	348.6	I	486.9	I	613.8
		II	486.1	II	609.1
		III	485.7	III	606.1
		IV	489.7	IV	609.0
Lanao del Sur	273.2	I	271.6	I	384.2
		II	263.7	II	341.7
		III	260.0	III	314.0
		IV	304.7	IV	340.3
Maguindanao	181.2	I	145.0	I	265.8
		II	137.5	II	220.8
		III	134.1	III	193.8
		IV	179.7	IV	219.4
Sulu	289.0	I	198.8	I	248.5
		II	188.0	II	196.9
		III	183.0	III	168.4
		IV	250.7	IV	195.4
Tawi-Tawi	298.0	I	118.6	I	272.5
		II	108.0	II	203.1
		III	103.3	III	164.8
		IV	168.8	IV	201.1

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of August

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening  
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity

CCI Category:  
■ Inadequate  
■ Sufficient  
■ Excess

## Regional Summary



The provinces of Basilan and Lanao del Sur received excessive rainfall amount for both rice and corn crops at all cropping stages. Such amount of rainfall may influence the expected production of corn crops especially those at the prevailing reproductive stage.

Based on SPEI3, majority of the region had near-normal to slightly wetter condition during the past 3 months. This signifies that moisture supply was available for both crops in most parts of the region. The Rx1day (1-day maximum rainfall) varies from 50-100 mm, while Rx5day (5-day maximum rainfall) reaches to about 150-200 mm. Based on the Rx5day, the magnitude of the extreme rainfall in the northern region was higher as compared to those in the southern part (Basilan, Sulu, and Tawi-tawi).

## NDVI

NDVI satellite data not yet available.

## SPEI3 (Jun-Jul-Aug)



## RX1day



## RX5day

