



Region VI (Western Visayas)

Water Availability for Rice
Prevailing Stage : (II) Vegetative



Water Availability for Corn
Prevailing Stage : (I) Establishment



Provincial Values

| Province | Mon. Ave. Rainfall (mm) | Rice | | | | Corn | | | |
|-------------------|-------------------------|------|---------|-----|---------|------|---------|-----|---------|
| | | CS | CCI (%) | CS | CCI (%) | CS | CCI (%) | CS | CCI (%) |
| Aklan | 384.1 | I | 232.7 | I | 423.6 | I | 89.5 | I | 152.5 |
| | | II | 219.4 | II | 336.7 | II | 85.1 | II | 131.9 |
| | | III | 213.4 | III | 285.2 | III | 82.9 | III | 117.4 |
| | | IV | 292.7 | IV | 334.1 | IV | 107.3 | IV | 131.2 |
| Antique | 238.9 | I | 169.7 | I | 333.3 | I | 119.5 | I | 246.7 |
| | | II | 156.8 | II | 246.9 | II | 109.2 | II | 174.8 |
| | | III | 150.8 | III | 197.8 | III | 104.6 | III | 135.6 |
| | | IV | 229.7 | IV | 244.3 | IV | 168.5 | IV | 172.7 |
| Capiz | 284.5 | I | 133.0 | I | 249.4 | I | 171.4 | I | 326.2 |
| | | II | 123.6 | II | 187.3 | II | 160.0 | II | 246.4 |
| | | III | 119.2 | III | 152.5 | III | 154.6 | III | 202.5 |
| | | IV | 174.7 | IV | 185.5 | IV | 223.6 | IV | 244.1 |
| Guimaras | 211.1 | I | 119.5 | I | 246.7 | I | 171.4 | I | 326.2 |
| | | II | 109.2 | II | 174.8 | II | 160.0 | II | 246.4 |
| | | III | 104.6 | III | 135.6 | III | 154.6 | III | 202.5 |
| | | IV | 168.5 | IV | 172.7 | IV | 223.6 | IV | 244.1 |
| Iloilo | 210.5 | I | 171.4 | I | 326.2 | I | 171.4 | I | 326.2 |
| | | II | 160.0 | II | 246.4 | II | 160.0 | II | 246.4 |
| | | III | 154.6 | III | 202.5 | III | 154.6 | III | 202.5 |
| | | IV | 223.6 | IV | 244.1 | IV | 223.6 | IV | 244.1 |
| Negros Occidental | 261.3 | I | 171.4 | I | 326.2 | I | 171.4 | I | 326.2 |
| | | II | 160.0 | II | 246.4 | II | 160.0 | II | 246.4 |
| | | III | 154.6 | III | 202.5 | III | 154.6 | III | 202.5 |
| | | IV | 223.6 | IV | 244.1 | IV | 223.6 | IV | 244.1 |

Crop Stage (CS) highlighted in black as the dominant stage during the month of December



Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening
Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity
CCI Category: Inadequate (yellow), Sufficient (green), Excess (blue)

Regional Summary



Rainfall was sufficient for the rice crops in the prevailing vegetative stage in most of the provinces. Meanwhile, the same crops in Aklan were likely affected by excess rainfall. For the corn crops, only the province of Antique had sufficient rainfall for the Establishment stage. The rest of the provinces had excessive rainfall for the same crop, nevertheless, the moisture was sufficient for the other stages of corn over Antique, Guimaras and Iloilo.

Crops are suggested to be in good condition in most of the farm areas in Western Visayas (see NDVI). This is despite the consecutively dry months since October. Meanwhile, excess one- to five-day rainfall had affected the southern to western part of the Region (see RX1day and RX5day), which is likely due to TC influence, the shear line, or the ITCZ.


NDVI



SPEI3 (Oct-Nov-Dec)



RX1day



RX5day

