




# Region X (Northern Mindanao)



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bukidnon	346.0	I	257.1	I	445.1
		II	241.8	II	343.5
		III	234.9	III	285.8
		IV	327.4	IV	340.5
Camiguin	356.8	I	332.7	I	477.6
		II	315.6	II	377.5
		III	307.8	III	318.5
		IV	407.7	IV	374.5
Lanao del Norte	385.7	I	255.3	I	465.2
		II	239.6	II	356.7
		III	232.4	III	295.8
		IV	328.7	IV	353.5
Misamis Occidental	517.1	I	331.6	I	558.9
		II	313.9	II	455.9
		III	305.8	III	392.2
		IV	409.8	IV	452.7
Misamis Oriental	410.2	I	278.9	I	493.6
		II	262.0	II	375.8
		III	254.3	III	310.3
		IV	357.7	IV	372.3

Crop Stage (CS) highlighted in black is the dominant stage during the month of December



## Regional Summary

The region received excessive rainfall during the month which might be detrimental to rice crops especially during their nursery to vegetative stage. Meanwhile, land preparation is ongoing for the corn crops.

The SPEI3 indicates somewhat slightly dry to dry conditions over the last three months throughout the entire region, whereas the NDVI indicates healthy crop conditions for the month. The rainfall extremes indicate that there is a maximum 1-day rainfall of 100 to 150 millimeters and a maximum 5-day rainfall of 150 to 250 millimeters respectively. ITCZ, shearline, and localized thunderstorms could be the primary cause of the month's heavy rainfall, as well as the extreme rainfall event.

