


# Region III (Central Luzon)

Water Availability for Rice  
Prevailing Stage : (II) Vegetative




Water Availability for Corn  
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

## Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice CCI (%)				Corn CCI (%)			
		CS I	CS II	CS III	CS IV	CS I	CS II	CS III	CS IV
Aurora	99.3	I	44.6	I	87.1	I	41.4	II	65.8
		II	39.9	III	53.1	II	9.8	II	15.5
		III	59.1	IV	65.1	III	9.2	III	11.6
		IV	10.9	I	21.7	IV	15.5	IV	15.3
Bataan	37.4	I	30.0	I	55.8	I	15.0	I	30.6
		II	27.6	II	39.7	II	13.6	II	22.2
		III	26.5	III	30.9	III	13.0	III	17.4
		IV	41.4	IV	39.2	IV	21.2	IV	22.0
Bulacan	87.9	I	6.5	I	14.9	I	1.6	I	4.7
		II	5.8	II	9.9	II	1.4	II	2.9
		III	5.5	III	7.3	III	1.3	III	2.1
		IV	10.0	IV	9.8	IV	2.6	IV	2.9
Nueva Ecija	56.0	I	8.0	I	22.9	I	8.0	I	22.9
		II	7.0	II	14.3	II	7.0	II	14.3
		III	6.6	III	10.1	III	6.6	III	10.1
		IV	13.3	IV	14.1	IV	13.3	IV	14.1
Pampanga	31.3	I	1.6	I	4.7	I	1.6	I	4.7
		II	1.4	II	2.9	II	1.4	II	2.9
		III	1.3	III	2.1	III	1.3	III	2.1
		IV	2.6	IV	2.9	IV	2.6	IV	2.9
Tarlac	25.0	I	8.0	I	22.9	I	8.0	I	22.9
		II	7.0	II	14.3	II	7.0	II	14.3
		III	6.6	III	10.1	III	6.6	III	10.1
		IV	13.3	IV	14.1	IV	13.3	IV	14.1
Zambales	44.2	I	8.0	I	22.9	I	8.0	I	22.9
		II	7.0	II	14.3	II	7.0	II	14.3
		III	6.6	III	10.1	III	6.6	III	10.1
		IV	13.3	IV	14.1	IV	13.3	IV	14.1

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of February




## Regional Summary

The rainfall was inadequate for most of the farm areas in the Region, except for some crops in the non-prevailing stages in Aurora and Bulacan. Meanwhile, the SPEI3 suggests slightly above normal moisture conditions over most of Nueva Ecija, Tarlac, Zambales, Pampanga and Bataan. Both the 1-day and 5-day maximum consecutive rainfall also report minimal rainfall amounts (i.e., maximum of ~50mm to ~100 mm, respectively).

## NDVI

NDVI satellite data not yet available.

## SPEI3 (Dec-Jan-Feb)



## RX1day



## RX5day

