



Region X (Northern Mindanao)

Water Availability for Rice
Prevailing Stage : (III) Reproductive




Water Availability for Corn
Prevailing Stage : (I) Establishment



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bukidnon	268.0	I	213.8	I	360.1
		II	201.1	II	278.3
		III	195.3	III	231.6
		IV	271.1	IV	275.9
Camiguin	299.8	I	267.9	I	370.8
		II	252.7	II	285.7
		III	245.7	III	232.6
		IV	336.8	IV	283.2
Lanao del Norte	234.1	I	246.1	I	401.0
		II	237.5	II	343.2
		III	233.4	III	309.3
		IV	283.7	IV	341.4
Misamis Occidental	200.0	I	258.6	I	327.0
		II	254.8	II	309.5
		III	253.0	III	296.2
		IV	273.8	IV	308.9
Misamis Oriental	394.5	I	304.7	I	570.8
		II	286.8	II	441.9
		III	278.7	III	369.0
		IV	387.0	IV	438.1

Crop Stage (CS) highlighted in black is the dominant stage during the month of February



Regional Summary



Almost all of the provinces in the Region received excessive rainfall during the month which might be detrimental to the rice crops that are mostly in the Reproductive stage, except over Bukidnon where rainfall was sufficient. Corn crops in the Region also received excessive rainfall where they are predominantly in the Establishment stage.

Generally, near-normal conditions have prevailed in most parts of the region, with slightly below normal rainfall over most parts of Misamis Occidental and Bukidnon (see SPEI3). Moreover, RX1day suggests that no significant rainfall event occurred in the area, even if RX5day had a slightly higher rainfall amount over Misamis Oriental, which could have been attributed to the Intertropical Convergence Zone (ITCZ), easterlies, localized thunderstorms, and shear line.

NDVI

NDVI satellite data not yet available.

SPEI3 (Dec-Jan-Feb)



RX1day



RX5day

