



# Bangsamoro Autonomous Region in Muslim Mindanao



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Basilan	87.7	I	24.6	I	46.0
		II	22.7	II	32.9
		III	21.9	III	26.0
		IV	21.9	IV	32.6
Lanao del Sur	177.6	I	106.6	I	188.2
		II	99.8	II	146.5
		III	96.6	III	121.3
		IV	96.6	IV	145.2
Maguindanao	212.1	I	128.5	I	239.7
		II	119.6	II	178.2
		III	115.6	III	144.7
		IV	115.6	IV	176.4
Sulu	78.6	I	30.1	I	45.5
		II	27.4	II	31.2
		III	26.2	III	23.9
		IV	26.2	IV	30.8
Tawi-Tawi	77.3	I	16.4	I	49.1
		II	14.4	II	32.4
		III	13.5	III	23.0
		IV	13.5	IV	31.9

Crop Stage (CS) highlighted in black is the dominant stage during the month of January.

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening  
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity

CCI Category:  
 Inadequate (Yellow)  
 Sufficient (Green)  
 Excess (Blue)

## Regional Summary

Basilan, Sulu, and Tawi-Tawi received an inadequate amount of water supply during the month, which may have adverse effects on rice and corn crops in the prevailing ripening and reproductive stages, respectively. On the other hand, Lanao del Sur and Maguindanao received sufficient rainfall to support rice and corn crops in any cropping stages.

Meanwhile, the SPEI3 indicated that the region was generally in near-normal condition, except for Basilan, some portions of Sulu, and Tawi-Tawi, where slightly drier conditions were observed. The ranges given by the RX1day (maximum 1-day rainfall) and RX5day (maximum 5-day rainfall) indices are both around 50–100 mm, where no notable extreme rainfall events that might have caused harm on rice or corn crops in the region.

