


Bangsamoro Autonomous Region in Muslim Mindanao



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Basilan	193.5	I	116.3	I	197.1
		II	108.2	II	154.4
		III	104.2	III	128.6
		IV	104.2	IV	153.1
Lanao del Sur	299.7	I	246.6	I	377.3
		II	236.1	II	318.6
		III	231.2	III	281.8
		IV	231.2	IV	316.8
Maguindanao	184.3	I	278.2	I	378.0
		II	275.0	II	354.7
		III	273.6	III	340.3
		IV	273.6	IV	354.0
Sulu	186.4	I	140.7	I	201.8
		II	131.2	II	155.7
		III	126.8	III	127.5
		IV	126.8	IV	154.3
Tawi-Tawi	247.3	I	148.3	I	265.2
		II	139.4	II	205.9
		III	135.3	III	170.9
		IV	135.3	IV	204.1

Crop Stage (CS) highlighted in **black** is the dominant stage during the month of January.

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity

CCI Category:
 Inadequate
 Sufficient
 Excess

Regional Summary

Rainfall in Basilan, Sulu, and Tawi-Tawi was sufficient to support the prevailing ripening and reproductive stages of rice and corn, respectively. On the other hand, excessive rainfall was observed in Lanao del Sur and Maguindanao. Such excessive rainfall could cause damage for both rice and corn crops in the absence of an effective drainage system in the area. Despite the excessive rainfall in some provinces, the NDVI still shows a high vegetation value in the entire region.

For the past three months, the region has generally been in slightly wetter conditions, except in Lanao del Sur and some parts of Maguindanao. The wetter conditions and excessive rainfall in these areas may be attributed to the Easterlies and localized thunderstorms.

