



Region VI (Western Visayas)

Water Availability for Rice
Prevailing Stage : (III) Reproductive



Water Availability for Corn
Prevailing Stage : (II) Vegetative



Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice				Corn			
		CS	CCI (%)	CS	CCI (%)	CS	CCI (%)	CS	CCI (%)
Aklan	202.7	I	100.7	I	185.9	I	100.7	I	185.9
		II	93.7	II	141.9	II	93.7	II	141.9
		III	90.5	III	115.1	III	90.5	III	115.1
		IV	90.5	IV	140.5	IV	90.5	IV	140.5
Antique	111.3	I	28.1	I	48.2	I	28.1	I	48.2
		II	26.4	II	40.8	II	26.4	II	40.8
		III	25.7	III	35.5	III	25.7	III	35.5
		IV	25.7	IV	40.5	IV	25.7	IV	40.5
Capiz	134.1	I	60.5	I	132.0	I	60.5	I	132.0
		II	55.1	II	91.0	II	55.1	II	91.0
		III	52.7	III	69.3	III	52.7	III	69.3
		IV	52.7	IV	89.9	IV	52.7	IV	89.9
Guimaras	88.8	I	28.2	I	68.1	I	28.2	I	68.1
		II	25.6	II	46.0	II	25.6	II	46.0
		III	24.5	III	35.2	III	24.5	III	35.2
		IV	24.5	IV	45.4	IV	24.5	IV	45.4
Iloilo	98.5	I	37.8	I	80.4	I	37.8	I	80.4
		II	34.7	II	56.3	II	34.7	II	56.3
		III	33.4	III	44.1	III	33.4	III	44.1
		IV	33.4	IV	55.7	IV	33.4	IV	55.7
Negros Occidental	137.1	I	70.8	I	161.1	I	70.8	I	161.1
		II	64.4	II	113.8	II	64.4	II	113.8
		III	61.6	III	87.3	III	61.6	III	87.3
		IV	61.6	IV	112.4	IV	61.6	IV	112.4

Crop Stage (CS) highlighted in black to the dominant stage during the month of January


Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening
Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity
CCI Category: Inadequate (yellow), Sufficient (green), Excess (blue)

Regional Summary

Rainfall was sufficient for the rice crops in the prevailing Reproductive stage in Aklan, Capiz and Negros Occidental. Meanwhile, the development of the rice crops in Antique, Guimaras, and Iloilo were likely hampered by inadequate rainfall. For the corn crops, the provinces of Antique and Guimaras had inadequate rainfall for the Vegetative stage. The rest of the provinces had sufficient rainfall for corn crops.

Both crops are suggested to be in good condition in most of the farm areas in Western Visayas (see NDVI) likely due to the generally near normal moisture available in the past three months (see SPEI3). Meanwhile, excess one- to five-day rainfall had affected the northern part of the Region (see RX1day and RX5day), which is likely due to the shear line or the ITCZ.

NDVI



SPEI3 (Nov-Dec-Jan)



RX1day



RX5day

