




# Bangsamoro Autonomous Region in Muslim Mindanao



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Basilan	295.7	I	214.2	I	336.3
		II	204.7	II	277.9
		III	200.2	III	241.2
		IV	255.2	IV	276.0
Lanao del Sur	350.7	I	531.9	I	636.0
		II	531.6	II	633.6
		III	531.4	III	632.0
		IV	533.4	IV	633.6
Maguindanao	261.7	I	449.3	I	605.1
		II	447.0	II	589.0
		III	446.0	III	578.8
		IV	458.7	IV	588.5
Sulu	306.4	I	203.8	I	301.9
		II	191.2	II	228.6
		III	185.5	III	186.3
		IV	262.1	IV	226.4
Tawi-Tawi	285.4	I	189.0	I	318.8
		II	177.4	II	246.1
		III	172.2	III	204.8
		IV	241.5	IV	243.9

Crop Stage (CS) highlighted in black to the dominant stage during the month of July



## Regional Summary

Most provinces in the region received excessive rainfall during the month, which may have damaged the corn crops at the prevailing reproductive stage and rice crops at the prevailing ripening stage, if not harvested earlier.

Based on SPEI3, majority of the region had near-normal rainfall during the past 3 months, which signifies that moisture supply was available for both crops in most of the growing period. But Lanao del Sur and Maguindanao had below-normal rainfall despite the excessive amount in the current month, signifying that crops may have suffered from dry conditions.

The RX1day varies from 50-150 mm, while RX5day is 200-250 mm. This suggests that, in the absence of drainage systems, standing crops may have been subjected to flooding.

