



# Region X (Northern Mindanao)



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bukidnon	433.2	I	518.8	I	796.7
		II	505.0	II	717.1
		III	498.6	III	668.5
		IV	580.0	IV	714.7
Camiguin	450.9	I	982.5	I	1078.6
		II	982.5	II	1078.6
		III	982.5	III	1078.6
		IV	982.5	IV	1078.6
Lanao del Norte	385.0	I	738.2	I	941.8
		II	738.2	II	941.8
		III	738.2	III	941.8
		IV	738.2	IV	941.8
Misamis Occidental	390.5	I	578.6	I	680.2
		II	578.6	II	680.2
		III	578.6	III	680.2
		IV	578.6	IV	680.2
Misamis Oriental	491.6	I	662.1	I	924.7
		II	650.9	II	846.6
		III	645.8	III	802.7
		IV	712.9	IV	844.4

Crop Stage (CS) highlighted in black is the dominant stage during the month of July.

Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening  
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity  
 CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

## Regional Summary

The amount of rainfall received this month was excessive which may have negatively affected the rice crops at all cropping stage in Northern Mindanao. Additionally, the 3-month Standardized Precipitation Evapotranspiration Index (SPEI) map indicates slightly wetter conditions, with a notable area experiencing significantly wetter conditions in Misamis Oriental, while the remaining provinces in the region are experiencing near-normal to slightly wetter conditions.

Meanwhile, the extreme rainfall indices RX1day and RX5day show significant rainfall (50-100 mm and 150-300 mm, respectively), which could have caused damage to rice crops (primarily in the central portion of the region, particularly in Bukidnon) due to heavy rainfall caused by the combined effect of ITCZ, low-pressure area, southwest monsoon, and localized thunderstorms.

