



# Bangsamoro Autonomous Region in Muslim Mindanao

**Water Availability for Rice**  
Prevailing Stage : (III) Reproductive




**Water Availability for Corn**  
Prevailing Stage : (II) Vegetative



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice				Corn			
		CS	CCI (%)	CS	CCI (%)	CS	CCI (%)	CS	CCI (%)
Basilan	413.9	I	630.3	I	690.4	I	630.3	II	670.9
		II	630.3	II	670.9	II	630.3	III	658.9
		III	630.3	III	658.9	III	630.3	IV	670.3
		IV	630.3	IV	670.3	IV	630.3	I	309.2
Lanao del Sur	258.5	I	196.9	I	309.2	I	188.4	II	258.3
		II	188.4	II	258.3	II	188.4	III	227.8
		III	188.4	III	227.8	III	188.4	IV	235.1
		IV	235.1	IV	256.7	IV	235.1	I	458.9
Maguindanao	224.3	I	328.4	I	458.9	I	323.4	II	426.9
		II	323.4	II	426.9	II	323.4	III	407.4
		III	323.4	III	407.4	III	323.4	IV	425.9
		IV	425.9	IV	425.9	IV	323.4	I	334.1
Sulu	266.2	I	346.4	I	334.1	I	337.5	II	286.6
		II	337.5	II	286.6	II	337.5	III	257.2
		III	337.5	III	257.2	III	337.5	IV	285.1
		IV	285.1	IV	285.1	IV	193.0	I	371.7
Tawi-Tawi	347.8	I	193.0	I	371.7	I	182.6	II	284.3
		II	182.6	II	284.3	II	182.6	III	235.2
		III	182.6	III	235.2	III	182.6	IV	243.0
		IV	243.0	IV	281.8	IV	243.0	I	334.1

Crop Stage (CS) highlighted in black is the dominant stage during the month of June




## Regional Summary

The BARMM region received sufficient to excessive amount of rainfall during the month, which is advantageous for the growth of corn crops at the prevailing vegetative stage and rice crops at the prevailing reproductive stage.


According to the SPEI3 map, the majority of the region had near-normal rainfall during the past 3-months, but Lanao del Sur and Maguindanao had below-normal rainfall and significantly drier conditions.

The RX1day and RX5day indices show that the rainfall varies from 50 to 150 millimeters, supporting the crops during the month of their growth stages.


**NDVI**



**SPEI3 (Apr-May-Jun)**



**RX1day**



**RX5day**

