


Region XII (SOCCSKSARGEN)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : (II) Vegetative



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Cotabato	205.9	I	380.9	I	511.1
		II	379.4	II	503.4
		III	378.7	III	498.4
		IV	387.0	IV	503.1
Cotabato City	200.1	I	225.7	I	384.8
		II	218.5	II	340.6
		III	215.2	III	311.0
		IV	256.7	IV	339.2
Sarangani	199.1	I	227.8	I	287.5
		II	224.4	II	261.9
		III	222.9	III	246.5
		IV	242.8	IV	261.1
South Cotabato	194.2	I	102.5	I	152.5
		II	99.4	II	140.9
		III	97.9	III	132.8
		IV	114.6	IV	140.5
Sultan Kudarat	229.3	I	260.7	I	402.4
		II	253.5	II	359.6
		III	250.1	III	334.5
		IV	292.4	IV	358.3

Crop Stage (CS) highlighted in black to the dominant stage during the month of June



Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity
 CCI Category: Inadequate (yellow), Sufficient (green), Excess (blue)

Regional Summary



The majority of the provinces in the SOCCSKSARGEN region received excess water supply for rice and corn crops at all cropping stages, with the exception of South Cotabato, which received adequate rainfall to sustain rice and corn crops at all cropping stages.

Despite excessive water for this month, the 3-month SPEI map still captured slightly drier to near normal conditions in almost all parts of the region, except for some areas in South Cotabato where moderately drier conditions were observed. The maximum rainfall in the region, as seen in Rx1day and Rx5day maps, ranges from 50 mm to 100 mm.


NDVI



SPEI3 (Apr-May-Jun)



RX1day



RX5day



Cotabato Cotabato City Sarangani South Cotabato Sultan Kudarat

