



# Region V (Bicol Region)

Water Availability for Rice  
Prevailing Stage : (I) Nursery



Water Availability for Corn  
Prevailing Stage : (II) Vegetative



## Provincial Values

		Mon. Ave. Rainfall (mm)	Rice				Corn			
			CS	CCI (%)	CS	CCI (%)	CS	CCI (%)	CS	CCI (%)
Albay	267.7		I	146.1	I	250.2	I	146.1	I	250.2
			II	136.8	II	191.3	II	136.8	II	191.3
			III	132.5	III	157.9	III	132.5	III	157.9
			IV	189.3	IV	189.6	IV	189.3	IV	189.6
Camarines Norte	334.8		I	184.0	I	322.6	I	184.0	I	322.6
			II	172.7	II	247.9	II	172.7	II	247.9
			III	167.6	III	205.8	III	167.6	III	205.8
			IV	236.5	IV	245.7	IV	236.5	IV	245.7
Camarines Sur	360.4		I	209.4	I	397.0	I	209.4	I	397.0
			II	195.2	II	293.2	II	195.2	II	293.2
			III	188.7	III	238.2	III	188.7	III	238.2
			IV	278.6	IV	290.3	IV	278.6	IV	290.3
Catanduanes	325.4		I	184.7	I	313.2	I	184.7	I	313.2
			II	173.8	II	246.2	II	173.8	II	246.2
			III	168.8	III	207.1	III	168.8	III	207.1
			IV	234.8	IV	244.2	IV	234.8	IV	244.2
Masbate	358.8		I	222.5	I	418.3	I	222.5	I	418.3
			II	208.1	II	315.6	II	208.1	II	315.6
			III	201.6	III	260.0	III	201.6	III	260.0
			IV	290.5	IV	312.7	IV	290.5	IV	312.7
Sorsogon	289.4		I	177.3	I	338.0	I	177.3	I	338.0
			II	165.7	II	254.7	II	165.7	II	254.7
			III	160.5	III	209.8	III	160.5	III	209.8
			IV	232.3	IV	252.3	IV	232.3	IV	252.3

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of June

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening  
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity  
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Red)

## Regional Summary


Most of the provinces in the region experienced an excessive amount of rainfall, which could have potentially caused damage to rice crops at the prevailing Nursery stage in Camarines Sur and Masbate, and to corn crops at the prevailing Vegetative stage in all provinces except Albay. In contrast, Camarines Norte, Catanduanes, and Sorsogon received sufficient rainfall for rice crops, while Albay for both rice and corn crops.

The three-month accumulated rainfall (SPEI3) map indicates that the entire region has experienced near-normal to slightly wetter conditions over the past three months, particularly in the western section. The maximum rainfall indices (RX1day and RX5day) showed that rainfall reached between 100 and 250 mm, occurring primarily during the first and second weeks of the month. This significant rainfall may be attributed to a low-pressure area that affected the region.

## NDVI

NDVI satellite data not yet available.

## SPEI3 (Apr-May-Jun)



## RX1day



## RX5day

