



# Region VII (Central Visayas)

Water Availability for Rice  
Prevailing Stage : (II) Vegetative



Water Availability for Corn  
Prevailing Stage : (II) Vegetative



NDVI

NDVI satellite  
data not yet  
available.

## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bohol	340.0	I	225.4	I	419.3
		II	211.6	II	320.5
		III	205.3	III	266.2
		IV	290.5	IV	317.7
Cebu	346.4	I	398.6	I	588.1
		II	385.5	II	503.5
		III	379.4	III	453.6
		IV	456.6	IV	501.0
Negros Oriental	331.7	I	315.2	I	510.0
		II	302.4	II	420.8
		III	296.6	III	369.0
		IV	373.1	IV	418.2
Siquijor	451.5	I	411.8	I	627.8
		II	391.3	II	495.7
		III	381.9	III	419.7
		IV	503.1	IV	491.8

Crop Stage (CS) highlighted in black is the dominant stage during the month of June

Rice CS:  
(I) Nursery  
(II) Vegetative  
(III) Reproductive  
(IV) Ripening

Corn CS:  
(I) Establishment  
(II) Vegetative  
(III) Reproductive  
(IV) Maturity



CCI Category:  
Inadequate  
Sufficient  
Excess

## Regional Summary

The prevailing stage of rice and corn crops in the region is both in the Vegetative stage during the month. This stage signifies that both crops will need much water for their growth and development. However, due to the influence of the intertropical convergence zone (ITCZ), Southwest monsoon, and localized thunderstorms, the water supply in the region has exceeded the water requirements of both crops, potentially leading to water logging in the absence of a proper and effective drainage system in the area.

The abundance of water supply in the area is reflected on the SPEI3 map, indicating that the region is in wetter conditions. Additionally, the rainfall in the region was more pronounced during the first to third weeks of the month, with a maximum rainfall that was around 100-150 mm (see RX5day).

SPEI3 (Apr-May-Jun)



RX1day



RX5day

