

# Region VI (Western Visayas)

Water Availability for Rice  
Prevailing Stage : None

## Provincial Values

|                   | Mon. Ave.<br>Rainfall<br>(mm) | Rice |         | Corn |         |
|-------------------|-------------------------------|------|---------|------|---------|
|                   |                               | CS   | CCI (%) | CS   | CCI (%) |
| Aklan             | 143.8                         | I    | 63.0    | I    | 126.0   |
|                   |                               | II   | 57.9    | II   | 96.3    |
|                   |                               | III  | 55.5    | III  | 79.1    |
|                   |                               | IV   | 85.9    | IV   | 95.4    |
| Antique           | 128.9                         | I    | 29.8    | I    | 49.3    |
|                   |                               | II   | 27.7    | II   | 39.8    |
|                   |                               | III  | 26.7    | III  | 33.5    |
|                   |                               | IV   | 39.1    | IV   | 39.5    |
| Capiz             | 147.6                         | I    | 69.9    | I    | 148.6   |
|                   |                               | II   | 64.2    | II   | 105.2   |
|                   |                               | III  | 61.7    | III  | 82.8    |
|                   |                               | IV   | 97.7    | IV   | 104.0   |
| Guimaras          | 128.5                         | I    | 53.1    | I    | 121.4   |
|                   |                               | II   | 47.9    | II   | 85.4    |
|                   |                               | III  | 45.6    | III  | 65.4    |
|                   |                               | IV   | 78.1    | IV   | 84.3    |
| Iloilo            | 130.6                         | I    | 48.2    | I    | 103.4   |
|                   |                               | II   | 43.8    | II   | 70.1    |
|                   |                               | III  | 41.8    | III  | 53.0    |
|                   |                               | IV   | 70.6    | IV   | 69.2    |
| Negros Occidental | 154.2                         | I    | 68.1    | I    | 154.9   |
|                   |                               | II   | 61.8    | II   | 108.2   |
|                   |                               | III  | 59.0    | III  | 82.8    |
|                   |                               | IV   | 99.9    | IV   | 106.9   |

Crop Stage (CS) highlighted in black is the dominant stage during the month of March

Rice CS:

(I) Nursery

(II) Vegetative

(III) Reproductive

(IV) Ripening

Corn CS:

(I) Establishment

(II) Vegetative

(III) Reproductive

(IV) Maturity

CCI Category:

Inadequate

Sufficient

Excess

## Regional Summary


Both crops in Aklan, Capiz, Guimaras, Iloilo, and Negros Occidental received a sufficient amount of rainfall throughout the month. Meanwhile, in Antique, there was inadequate rainfall during the prevailing Maturity stage of corn crops.

The NDVI shows that crops are generally in healthy condition however, notable areas in Antique, Iloilo, Capiz and Negros Occidental are slightly stressed.



The SPEI3 suggests generally near normal conditions were experienced for the past three months in the region. Rainfall extremes show that the majority of the region received rainfall ranging from 50-150 mm, while 200mm maximum rainfall was observed in the northeast portion of Negros Occidental.

Harvesting and/or  
pre-planting  
stage is possibly  
ongoing.



Water Availability for Corn  
Prevailing Stage : (IV) Maturity



NDVI



SPEI3 (Jan-Feb-Mar)



RX1day



RX5day

