

Region IV-B (MIMAROPA)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Marinduque	298.3	I	76.3	I	198.8
		II	68.9	II	128.9
		III	65.7	III	94.7
		IV	117.8	IV	127.0
Occidental Mindoro	204.4	I	32.8	I	66.7
		II	29.9	II	43.3
		III	28.6	III	32.5
		IV	48.9	IV	42.7
Oriental Mindoro	201.8	I	53.2	I	115.3
		II	48.5	II	77.3
		III	46.4	III	58.9
		IV	78.0	IV	76.3
Palawan	109.3	I	20.9	I	50.0
		II	18.8	II	32.7
		III	17.9	III	24.3
		IV	31.7	IV	32.2
Romblon	296.5	I	64.4	I	139.4
		II	59.1	II	96.0
		III	56.7	III	74.5
		IV	91.8	IV	94.8

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of May

Rice CS:
(I) Nursery
(II) Vegetative
(III) Reproductive
(IV) Ripening

Corn CS:
(I) Establishment
(II) Vegetative
(III) Reproductive
(IV) Maturity

CCI Category:
Inadequate
Sufficient
Excess

Regional Summary

Throughout the month, most of the region's provinces received enough water to sustain rice and corn crops at all cropping stages, with the exception of Occidental Mindoro and Palawan. Likewise, the SPEI3 map indicated that the western section of the region had slightly drier to near normal conditions, while the eastern portions of the region experienced slightly wetter to near normal conditions. This can be seen in RX1day and RX5day maps, as rainfall appeared to be concentrated in the eastern portion of the region, which is roughly 100–300 mm. This rainfall is widely attributed to Typhoon AGHON, which passed through the eastern section of the region.

Despite the passage of the typhoon, there was no significant damage to rice and corn crops in the region, as there were no prevailing crops during that period.

