



Region VII (Central Visayas)

Water Availability for Rice
Prevailing Stage : (I) Nursery



Water Availability for Corn
Prevailing Stage : (I) Establishment



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bohol	57.3	I	7.2	I	18.0
		II	6.3	II	11.4
		III	5.9	III	7.9
		IV	12.0	IV	11.2
Cebu	92.5	I	14.6	I	33.3
		II	13.2	II	21.7
		III	12.6	III	16.3
		IV	21.7	IV	21.4
Negros Oriental	58.6	I	8.3	I	24.8
		II	7.1	II	13.9
		III	6.6	III	9.2
		IV	15.0	IV	13.7
Siquijor	73.9	I	12.5	I	31.6
		II	10.9	II	19.6
		III	10.2	III	14.0
		IV	20.4	IV	19.3

Crop Stage (CS) highlighted in black is the dominant stage during the month of May

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)


Regional Summary

All provinces in the region received an inadequate amount of rainfall throughout the month for both rice and corn in all cropping stages. This shortage could have possible adverse effects on rice crops in the nursery stage and affect the growth and development of corn crops. The scarcity of water is due to the limits of monthly rainfall, as seen in RX1day and RX5day.

Additionally, the region experienced a slightly dry condition, except in some areas in Bohol where there were significantly drier conditions, as shown in SPEI 3. It is also evident that most of the crops in the farm areas in the region were in stressed condition.

No notable extreme rainfall events affected the region during the month. The extreme rainfall indices showed maximum 1-day and 5-day rainfall estimates of 50mm-150mm in portions of Cebu and Bohol.

NDVI



SPEI3 (Mar-Apr-May)



RX1day



RX5day

