



Region VI (Western Visayas)

Water Availability for Rice
Prevailing Stage : (I) Nursery



Water Availability for Corn
Prevailing Stage : (I) Establishment



Provincial Values

| | Mon. Ave. Rainfall (mm) | Rice | | | | Corn | | | |
|-------------------|-------------------------------|------|---------|-----|---------|------|---------|-----|---------|
| | | CS | CCI (%) | CS | CCI (%) | CS | CCI (%) | CS | CCI (%) |
| Aklan | 355.1 | I | 212.6 | I | 385.0 | I | 199.6 | II | 302.7 |
| | | II | 199.6 | II | 302.7 | II | 193.5 | III | 254.5 |
| | | III | 193.5 | III | 254.5 | III | 270.4 | IV | 300.3 |
| | | IV | 270.4 | IV | 300.3 | IV | 123.9 | I | 172.4 |
| Antique | 305.0 | I | 123.9 | I | 172.4 | I | 117.4 | II | 146.9 |
| | | II | 117.4 | II | 146.9 | II | 114.2 | III | 129.1 |
| | | III | 114.2 | III | 129.1 | III | 149.7 | IV | 146.0 |
| | | IV | 149.7 | IV | 146.0 | IV | 121.2 | I | 255.2 |
| Capiz | 265.8 | I | 121.2 | I | 255.2 | I | 111.5 | II | 177.8 |
| | | II | 111.5 | II | 177.8 | II | 107.3 | III | 139.1 |
| | | III | 107.3 | III | 139.1 | III | 170.2 | IV | 175.7 |
| | | IV | 170.2 | IV | 175.7 | IV | 213.4 | I | 393.4 |
| Guimaras | 375.0 | I | 213.4 | I | 393.4 | I | 198.1 | II | 288.3 |
| | | II | 198.1 | II | 288.3 | II | 191.2 | III | 228.4 |
| | | III | 191.2 | III | 228.4 | III | 153.0 | I | 285.6 |
| | | IV | 286.0 | IV | 285.1 | IV | 141.2 | II | 204.2 |
| Iloilo | 293.0 | I | 153.0 | I | 285.6 | I | 135.9 | III | 160.8 |
| | | II | 141.2 | II | 204.2 | II | 211.7 | IV | 201.9 |
| | | III | 135.9 | III | 160.8 | III | 131.4 | I | 250.0 |
| | | IV | 211.7 | IV | 201.9 | IV | 120.8 | II | 179.1 |
| Negros Occidental | 247.5 | I | 131.4 | I | 250.0 | I | 116.0 | III | 141.2 |
| | | II | 120.8 | II | 179.1 | II | 181.4 | IV | 177.0 |
| | | III | 116.0 | III | 141.2 | III | 181.4 | IV | 177.0 |
| | | IV | 181.4 | IV | 177.0 | IV | 181.4 | IV | 177.0 |

Crop Stage (CS) highlighted in black is the dominant stage during the month of May

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

Regional Summary



During the month, the prevailing stage of rice and corn crops is both in its early stage. For the prevailing nursery stage of rice crops, the region had a sufficient water supply to sustain its growth and development, except in Guimaras and Aklan, where the water supply was excessive. On the other hand, the prevailing establishment stage of corn crops in the region had an excessive water supply for the month, except in Antique, where a sufficient water supply was observed.

Those excessive waters in some areas of the region may be attributed to the ITCZ, LPA, and localized thunderstorms, which brought a maximum 5-day rainfall ranging from 100 to 250 mm. Consequently, the SPEI3 reflected a positive value, showing that the region was generally in wetter condition (except in the central part of Panay Island). Thus, proper drainage is advisable in those areas to attain good crop production and yield.

NDVI

NDVI satellite data not yet available.

SPEI3 (Mar-Apr-May)



RX1day



RX5day

