



# Region VII (Central Visayas)

Water Availability for Rice  
Prevailing Stage : (I) Nursery



Water Availability for Corn  
Prevailing Stage : (IV) Maturity



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bohol	104.9	I	41.0	I	90.8
		II	37.0	II	60.2
		III	35.2	III	44.5
		IV	35.2	IV	59.3
Cebu	98.6	I	45.0	I	95.3
		II	40.5	II	60.6
		III	38.5	III	44.4
		IV	38.5	IV	59.7
Negros Oriental	62.3	I	18.8	I	50.8
		II	16.4	II	30.4
		III	15.3	III	20.8
		IV	15.3	IV	29.9
Siquijor	208.0	I	122.4	I	226.9
		II	113.0	II	167.3
		III	108.6	III	134.4
		IV	108.6	IV	165.6

Crop Stage (CS) highlighted in black is the dominant stage during the month of November


Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening  
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity  
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

## Regional Summary


The Region's rainfall was sufficient for the following: corn crops at the prevailing maturity stage in the provinces of Bohol, Cebu, and Siquijor; rice crops at the prevailing nursery stage in province of Siquijor. Meanwhile, the provinces of Bohol, Cebu and Siquijor received inadequate amount of rainfall for their rice crops, this amount may affect the expected crop production for the month of November.

The three-month accumulated rainfall (SPEI3) indicates generally near-normal with slightly wetter conditions on the province of Siquijor and northern part of Cebu and Negros Oriental. Based on the RX5day, the maximum 5-day consecutive rainfall ranges from 50 to 150 mm indicating that no significant extreme rainfall that could have had a negative impact on crops in the area was observed. The NDVI map shows that the vegetation is in healthy condition especially those with NDVI values of greater than 0.8.

NDVI



SPEI3 (Sep-Oct-Nov)



RX1day



RX5day

