



# Region XI (Davao Region)

Water Availability for Rice  
Prevailing Stage : (I) Nursery



Water Availability for Corn  
Prevailing Stage : None



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS CCI (%)	CS CCI (%)	CS CCI (%)	CS CCI (%)
Compostela Valley	235.3	I	109.6	I	222.5
		II	99.8	II	162.1
		III	95.5	III	126.4
		IV	95.5	IV	160.2
Davao del Norte	177.3	I	77.2	I	144.9
		II	71.0	II	104.1
		III	68.2	III	82.5
		IV	68.2	IV	103.0
Davao del Sur	130.1	I	43.5	I	75.1
		II	39.7	II	53.6
		III	38.0	III	41.9
		IV	38.0	IV	53.0
Davao Occidental	157.7	I	42.9	I	86.6
		II	39.4	II	60.5
		III	37.9	III	47.4
		IV	37.9	IV	59.8
Davao Oriental	265.9	I	96.6	I	211.2
		II	87.6	II	148.8
		III	83.6	III	115.0
		IV	83.6	IV	147.0

Crop Stage (CS) highlighted in **black** is the dominant stage during the month of November



Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening  
 Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity  
 CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

## Regional Summary



Rice crops were mostly in the nursery stage, while corn crops were in the harvesting or pre-planting stages. The rainfall received for the month was mostly sufficient for all crops across the region except Davao del Sur and Davao Occidental, which received inadequate water.

The SPEI3 shows slightly drier conditions experienced for the past three months in the whole region with near healthy crop conditions in some parts spread across the region, indicated in the NDVI map. Additionally, the rainfall extremes indicate that the maximum 1-day rainfall estimated 50-100 mm but the maximum 5-day rainfall estimated amounts up to 200 mm.


NDVI



SPEI3 (Sep-Oct-Nov)



RX1day



RX5day

