


# Region II (Cagayan Valley)



## Provincial Values

	Mon. Ave. Rainfall (mm)	Rice CS CCI (%)	Corn CS CCI (%)
Batanes	298.0	I 82.3	I 165.2
		II 75.8	II 118.4
		III 72.9	III 93.5
		IV 72.9	IV 117.1
Cagayan	119.2	I 41.3	I 91.1
		II 37.1	II 62.0
		III 35.3	III 46.6
		IV 35.3	IV 61.2
Isabela	131.6	I 38.7	I 82.5
		II 35.3	II 56.9
		III 33.8	III 43.7
		IV 33.8	IV 56.1
Nueva Vizcaya	103.1	I 49.3	I 97.2
		II 45.1	II 69.5
		III 43.2	III 54.2
		IV 43.2	IV 66.7
Quirino	122.3	I 47.0	I 96.6
		II 43.0	II 66.0
		III 41.2	III 50.7
		IV 41.2	IV 65.2

## Regional Summary

The rainfall received during the month of October was sufficient for corn crops, especially those at the prevailing establishment stage in the region. However, the rainfall amount received for rice crops was inadequate for most provinces in Cagayan Valley, which may have affected the pre-planting activities in the region, except for Batanes. The NDVI map indicated generally fair condition which may also represent the establishment stage for the corn crops over the region (with NDVI value ranging from 0.5 to 0.75).

The SPEI3 (three-month accumulated rainfall) showed that most parts of the region experienced relatively drier conditions, except for major parts of Nueva Vizcaya province where near-normal condition was observed. The maximum 1-day (RX1day) and 5-day (RX5day) rainfall ranged from 50-100mm and 50-150mm, respectively.

