

Region V (Bicol Region)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : (I) Establishment



Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Albay	701.8	I	387.9	I	712.7
		II	356.3	II	549.1
		III	341.9	III	446.8
		IV	341.9	IV	544.1
Camarines Norte	595.5	I	336.6	I	577.4
		II	314.2	II	441.8
		III	303.9	III	360.0
		IV	303.9	IV	437.6
Camarines Sur	599.1	I	404.5	I	748.5
		II	373.9	II	563.1
		III	359.9	III	453.2
		IV	359.9	IV	557.5
Catanduanes	553.3	I	279.0	I	528.0
		II	255.5	II	403.6
		III	244.8	III	324.8
		IV	244.8	IV	399.7
Masbate	561.6	I	295.2	I	617.4
		II	267.6	II	426.6
		III	255.2	III	325.4
		IV	255.2	IV	421.1
Sorsogon	669.1	I	418.3	I	792.0
		II	382.8	II	597.5
		III	366.6	III	478.1
		IV	366.6	IV	591.6

Crop Stage (CS) highlighted in bold is the dominant stage during the month of October

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)

Regional Summary


During the month, the farm areas in the region suffered an excessive amount of water supply brought by the severe tropical storm Kristine. Maximum rainfall of about 400 mm was experienced during the fourth week of the month, which caused severe flooding in the area. According to the NDRRMC and DA-DRRM Operation Center for the combined effects of TC Kristine and Leon, the region received the highest production loss for agriculture in the country.

Furthermore, the region's SPEI for the previous three months shows that the majority of the area is in wetter conditions.


NDVI

NDVI satellite data not yet available.

SPEI3 (Aug-Sep-Oct)



RX1day



RX5day

