



# Region IX (Zamboanga Peninsula)

Water Availability for Rice  
Prevailing Stage : (IV) Ripening



Water Availability for Corn  
Prevailing Stage : (II) Vegetative



## Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
City of Isabela	245.7	I	80.2	I	162.2
		II	73.1	II	113.3
		III	69.9	III	88.2
		IV	69.9	IV	112.0
Zamboanga del Norte	290.2	I	206.2	I	318.0
		II	197.4	II	255.5
		III	193.4	III	218.3
		IV	193.4	IV	253.5
Zamboanga del Sur	233.0	I	156.6	I	249.4
		II	148.7	II	201.9
		III	145.1	III	173.1
		IV	145.1	IV	200.4
Zamboanga Sibugay	302.8	I	219.9	I	339.5
		II	211.0	II	286.3
		III	206.9	III	254.1
		IV	206.9	IV	284.6

Crop Stage (CS) highlighted in black is the dominant stage during the month of September


Rice CS: (I) Nursery, (II) Vegetative, (III) Reproductive, (IV) Ripening  
Corn CS: (I) Establishment, (II) Vegetative, (III) Reproductive, (IV) Maturity  
CCI Category: Inadequate (yellow), Sufficient (green), Excess (blue)

## Regional Summary



Rainfall had been excessive for the corn crops in most of the provinces (except in the City of Isabela), particularly for those in the prevailing Vegetative stage. Meanwhile, the rice crops in the ripening stage likely benefited from the sufficient rainfall, although certain areas had also experienced slightly higher volumes of rainfall.

The lower portion of the region has been having slightly higher than normal rainfall in the past 3 months, which, in the presence of an irrigation facility, is an opportunity to replenish its supply. Meanwhile, a notable accumulation of consecutive rainfall was found in the western portion of Zamboanga del Norte, which is possibly from the compounded effects of TS Ferdie and TD Gener.

NDVI




SPEI3 (Jul-Aug-Sep)



RX1day



RX5day



City of Isabela (green), Zamboanga del Norte (green), Zamboanga del Sur (red), Zamboanga Sibugay (red)

