


Region X (Northern Mindanao)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : (III) Reproductive



Provincial Values

Province	Mon. Ave. Rainfall (mm)	Rice CCI (%)		Corn CCI (%)	
		CS	CCI (%)	CS	CCI (%)
Bukidnon	68.3	I	2.5	I	14.3
		II	1.8	II	7.6
		III	1.6	III	3.4
		IV	8.3	IV	7.4
Camiguin	13.7	I	0.0	I	0.0
		II	0.0	II	0.0
		III	0.0	III	0.0
		IV	0.0	IV	0.0
Lanao del Norte	61.1	I	0.2	I	7.0
		II	0.1	II	0.5
		III	0.1	III	0.3
		IV	1.1	IV	0.5
Misamis Occidental	51.6	I	0.2	I	0.7
		II	0.2	II	0.4
		III	0.2	III	0.3
		IV	0.4	IV	0.4
Misamis Oriental	47.1	I	2.6	I	16.8
		II	1.6	II	10.7
		III	0.7	III	8.6
		IV	4.8	IV	10.6

Crop Stage (CS) highlighted in black to the dominant stage during the month of April

Rice CS:
(I) Nursery
(II) Vegetative
(III) Reproductive
(IV) Ripening

Corn CS:
(I) Establishment
(II) Vegetative
(III) Reproductive
(IV) Maturity



CCI Category:
 Inadequate
 Sufficient
 Excess

Regional Summary



The rainfall for the month was inadequate; this could have potentially caused damage on both rice and corn crops at any crop stage in Northern Mindanao. Moreover, the SPEI3 indicates slightly drier than normal condition in most parts of the region and generally near normal condition for the rest of the region.

There was no extreme rainfall that might have caused impact on both rice and corn crops in the entire region.


NDVI



SPEI3 (Feb-Mar-Apr)



RX1day



RX5day

