



# Cordillera Administrative Region

Water Availability for Rice  
Prevailing Stage : None



Water Availability for Corn  
Prevailing Stage : (III) Silking




## Provincial Values Regional Summary

	Mon. Ave. Rainfall (mm)	Rice				Corn			
		CS	RS (%)	CS	RS (%)	CS	RS (%)	CS	RS (%)
Mountain Province	23.68	I	29.0	I	39.7	I	29.0	I	39.7
		II	27.4	II	32.8	II	27.4	II	32.8
		III	26.7	III	28.4	III	26.7	III	28.4
		IV	26.7	IV	32.6	IV	26.7	IV	32.6
Kalinga	52.45	I	54.5	I	73.7	I	54.5	I	73.7
		II	51.8	II	61.7	II	51.8	II	61.7
		III	50.5	III	54.0	III	50.5	III	54.0
		IV	50.5	IV	61.4	IV	50.5	IV	61.4
Ifugao	26.28	I	27.7	I	38.2	I	27.7	I	38.2
		II	26.5	II	32.8	II	26.5	II	32.8
		III	26.0	III	29.2	III	26.0	III	29.2
		IV	26.0	IV	32.6	IV	26.0	IV	32.6
Benguet	24.08	I	23.2	I	34.9	I	23.2	I	34.9
		II	22.1	II	29.6	II	22.1	II	29.6
		III	21.6	III	26.1	III	21.6	III	26.1
		IV	21.6	IV	29.4	IV	21.6	IV	29.4
Apayao	78.34	I	81.1	I	141.9	I	81.1	I	141.9
		II	77.1	II	118.8	II	77.1	II	118.8
		III	75.3	III	103.9	III	75.3	III	103.9
		IV	75.3	IV	118.1	IV	75.3	IV	118.1
Abra	16.12	I	10.9	I	15.3	I	10.9	I	15.3
		II	10.3	II	12.4	II	10.3	II	12.4
		III	10.0	III	10.6	III	10.0	III	10.6
		IV	10.0	IV	12.3	IV	10.0	IV	12.3


Legend :  
Inadequate  
Sufficient  
Excess

Majority of the provinces in Cordillera Administrative region received insufficient amount of rainfall during the month of December. Only the provinces of Kalinga and Apayao received sufficient amount of rainfall for both rice and corn crops for all cropping stages.

Although healthy vegetation (NDVI > 0.25) was observed in most of the provinces, according to the 3-month accumulated rainfall, lower rainfall amount or drier condition was still observed in the southern part of the region. Furthermore, maximum 1-day and 5-day rainfall shows that there were no significant or extreme rainfall observed during this month.



## SPEI3 (Oct-Nov-Dec)



## RX1day



## RX5day

