


Region V (Bicol Region)



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	RS (%)	CS	RS (%)
Sorsogon	327.98	I	271.2	I	434.5
		II	253.2	II	334.3
		III	245.1	III	277.7
		IV	245.1	IV	331.4
Masbate	171.25	I	134.6	I	237.7
		II	125.5	II	181.9
		III	121.3	III	150.7
		IV	121.3	IV	180.3
Catanduanes	421.85	I	323.7	I	482.7
		II	304.9	II	392.2
		III	296.3	III	336.5
		IV	296.3	IV	389.4
Camarines Sur	320.31	I	213.0	I	381.6
		II	198.4	II	291.4
		III	191.8	III	241.0
		IV	191.8	IV	288.8
Camarines Norte	361.14	I	272.3	I	413.1
		II	256.1	II	331.4
		III	248.7	III	282.1
		IV	248.7	IV	328.9
Albay	262.33	I	181.4	I	249.5
		II	170.4	II	198.3
		III	165.3	III	167.8
		IV	165.3	IV	196.7

Legend :
Inadequate
Sufficient
Excess

Regional Summary

Most provinces in the Bicol region received excessive rainfall, which may have caused potential damage to both rice and corn crops in the prevailing nursery and silking stages, respectively. On the other hand, NDVI generally suggests a healthy vegetation in the region, with slightly low NDVI values over few parts of Camarines Norte and Camarines Sur.

Based on the SPEI3, accumulated rainfall for the past three months is generally near normal for most of the provinces, with slightly below normal conditions in the southern part of Sorsogon.

Due to the combined effects of shear line and LPA, notable heavy rainfall events have affected the provinces in the northern parts of the region as indicated in the RX1day and RX5day, with maximum accumulated rainfall amounting to 100mm and 300mm, respectively.

