



Region X (Northern Mindanao)



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	RS (%)	CS	RS (%)
Misamis Oriental	314.73	I	300.1	I	420.6
		II	283.2	II	340.5
		III	275.5	III	292.9
		IV	275.5	IV	338.1
Misamis Occidental	310.26	I	217.9	I	335.1
		II	205.0	II	269.3
		III	199.1	III	229.5
		IV	199.1	IV	267.3
Lanao del Norte	206.54	I	221.0	I	311.4
		II	209.9	II	256.1
		III	204.9	III	223.6
		IV	204.9	IV	254.4
Camiguin	486.86	I	522.5	I	894.1
		II	491.6	II	789.1
		III	477.5	III	714.9
		IV	477.5	IV	785.6
Bukidnon	263.12	I	278.9	I	403.9
		II	264.1	II	332.1
		III	257.4	III	289.0
		IV	257.4	IV	329.9

Legend :

- Inadequate
- Sufficient
- Excess

Regional Summary

Sufficient amount of rainfall received by the region during the month is beneficial in the preparation of rice paddy and also for the newly planted rice crops. NDVI suggests healthy vegetation across the region.

Based on the three-month consecutive rainfall as shown in the SPEI3, most of the provinces in Northern Mindanao and most parts of Bukidnon received near-normal to above normal amount of rainfall while in southern parts of Misamis Occidental, Lanao del Norte, and Southwest of Bukidnon, slightly below normal amount of rainfall was recorded.

Significantly heavy rainfall events indicated in the RX5day were experienced over the region in the latter part of the month. Rice crops in nursery stage could have been potentially damaged in some areas including the Misamis provinces and eastern part of Bukidnon.

