



Region XI (Davao Region)

Water Availability for Rice
Prevailing Stage : (II) Vegetative



Water Availability for Corn
Prevailing Stage : None



Provincial Values Regional Summary

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	RS (%)	CS	RS (%)
Davao Oriental	322.70	I	236.2	I	312.5
		II	221.3	II	247.8
		III	214.6	III	209.5
		IV	214.6	IV	245.8
Davao Occidental	191.99	I	126.5	I	191.0
		II	118.7	II	152.0
		III	115.2	III	128.9
		IV	115.2	IV	150.9
Davao del Sur	149.42	I	106.7	I	149.2
		II	100.2	II	119.1
		III	97.3	III	101.1
		IV	97.3	IV	118.2
Davao del Norte	268.41	I	138.0	I	231.6
		II	128.9	II	181.4
		III	124.8	III	152.4
		IV	124.8	IV	179.9
Compostela Valley	324.10	I	239.2	I	336.4
		II	224.9	II	269.9
		III	218.3	III	229.9
		IV	218.3	IV	267.9

Legend :
Inadequate
Sufficient
Excess

Rainfall for the month was sufficient to support rice crops at any stage in the western portion of the region. However, excessive rainfall may have caused damage to the rice crops in the eastern part of the region. Meanwhile, the NDVI suggests healthy vegetation in the farm areas of Davao region. The accumulated rainfall for the past three months was within the normal values for some areas, but slightly below normal rainfall was observed in the northeastern part of Davao del Sur and southern portion of Davao del Norte. based on the SPEI3.

Maximum 5-day rainfall of about 200-300mm was experienced at the northernmost portion of the region. These extreme events, which were mainly attributed to the Shearline and the presence of LPA may have caused an adverse impact on crops in the above-mentioned areas.

