

Region IV-B (MIMAROPA)

Water Availability for Rice
Prevailing Stage : (IV) Ripening



Water Availability for Corn
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Marinduque	16.9	I	0.1	I	0.5
		II	0.1	II	0.3
		III	0.1	III	0.2
		IV	0.2	IV	0.3
Occidental Mindoro	8.6	I	0.0	I	0.0
		II	0.0	II	0.0
		III	0.0	III	0.0
		IV	0.0	IV	0.0
Oriental Mindoro	23.0	I	0.9	I	1.9
		II	0.8	II	1.2
		III	0.7	III	0.9
		IV	1.4	IV	1.2
Palawan	11.8	I	0.1	I	0.3
		II	0.1	II	0.2
		III	0.1	III	0.1
		IV	0.2	IV	0.2
Romblon	13.8	I	0.3	I	0.4
		II	0.2	II	0.3
		III	0.2	III	0.2
		IV	0.4	IV	0.3

Crop Stage (CS) highlighted in **bold** is the dominant stage during the month of March

Rice CS:
(I) Nursery
(II) Vegetative
(III) Reproductive
(IV) Ripening

Corn CS:
(I) Establishment
(II) Vegetative
(III) Reproductive
(IV) Maturity

CCI Category:
 Inadequate
 Sufficient
 Excess

Regional Summary

The rainfall received for the month was inadequate. This could have caused potential damage on both rice and corn crops at any of the crop stages in the whole MIMAROPA region.

The NDVI suggest slightly stressed crops in some parts of Occidental Mindoro and Oriental Mindoro, with generally healthy conditions for the rest of the region. Moreover, the 3-month SPEI indicates generally near-normal conditions for the past three months in most parts of the region.

There was no significant extreme rainfall for the month that might have affected the rice and corn crops in the region, based on the RX1day and RX5day.

