


Region VII (Central Visayas)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or pre-planting stage is possibly ongoing.

Water Availability for Corn
Prevailing Stage : (IV) Maturity



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bohol	64.0	I	11.2	I	24.2
		II	10.1	II	16.4
		III	9.6	III	11.7
		IV	16.9	IV	16.2
Cebu	54.9	I	3.0	I	17.8
		II	2.3	II	8.3
		III	2.0	III	5.1
		IV	7.5	IV	8.0
Negros Oriental	40.3	I	0.7	I	7.9
		II	0.6	II	1.7
		III	0.5	III	0.4
		IV	3.3	IV	1.4
Siquijor	53.2	I	0.7	I	14.2
		II	0.6	II	7.5
		III	0.6	III	1.6
		IV	6.4	IV	7.1

Crop Stage (CS) highlighted in black is the dominant stage during the month of March

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening
Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity
CCI Category: Inadequate (Yellow), Sufficient (Green), Excess (Blue)


Regional Summary

In March, rainfall was inadequate in Central Visayas region. This may be harmful for growing crops, however, it is advantageous for corn crops reaching maturity. Additionally, it helps rice crops when they are ready for harvest.


The NDVI shows that existing are crops in generally good condition. The province of Bohol and certain areas of the region have near-normal to slightly drier weather, according to the 3-month cumulative rainfall index (SPEI3).

The trough of a Low Pressure Area (LPA) is likely responsible for the highest 1-day (RX1day) and maximum 5-day (RX5day) values of roughly 50–100 mm, respectively.

NDVI



SPEI3 (Jan-Feb-Mar)



RX1day



RX5day

