



Region XI (Davao Region)



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice CS	Rice CCI (%)	Corn CS	Corn CCI (%)
Compostela Valley	149.4	I	63.8	I	134.2
		II	58.2	II	98.1
		III	55.7	III	77.5
		IV	55.7	IV	97.0
Davao del Norte	231.3	I	115.7	I	208.3
		II	106.5	II	157.3
		III	102.4	III	127.0
		IV	102.4	IV	155.7
Davao del Sur	232.1	I	107.2	I	178.9
		II	98.9	II	132.6
		III	95.1	III	106.0
		IV	95.1	IV	131.3
Davao Occidental	67.6	I	15.6	I	41.6
		II	13.7	II	25.3
		III	12.9	III	17.7
		IV	12.9	IV	24.9
Davao Oriental	151.6	I	53.7	I	140.7
		II	48.5	II	101.4
		III	46.1	III	79.3
		IV	46.1	IV	100.3

Crop Stage (CS) highlighted in **blue** is the dominant stage during the month of October.

Rice CS: (I) Nursery (II) Vegetative (III) Reproductive (IV) Ripening

Corn CS: (I) Establishment (II) Vegetative (III) Reproductive (IV) Maturity

CCI Category: Inadequate Sufficient Excess

Regional Summary

In the provinces of Compostela Valley, Davao del Norte, and Davao del Sur, sufficient rainfall was observed for both rice and corn crops. This could be advantageous for the farmers in terms of preparing their land for crop planting. On the other hand, inadequate rainfall was received over the provinces of Davao Occidental and rice crops in Davao Oriental.

Meanwhile, the NDVI indicated that majority of the farm areas in the Davao Region seem to have healthy vegetation. In contrast, the SPEI3 indicated near normal to slightly drier conditions over some parts of Davao del Sur, Davao Oriental, and Davao Occidental. Moreover, maximum 1-day rainfall of 50-100mm and 5-day rainfall of 150-200mm were experienced over some parts of Davao del Norte, Davao del Sur, portions of Compostela Valley and Davao Oriental due to the combined effects southwest monsoon and localized thunderstorms, which may have detrimental effects to the farm areas in the region.

