


Region VII (Central Visayas)

Water Availability for Rice
Prevailing Stage : None

Harvesting and/or
pre-planting
stage is possibly
ongoing.

Water Availability for Corn
Prevailing Stage : (II) Vegetative



Provincial Values

	Mon. Ave. Rainfall (mm)	Rice		Corn	
		CS	CCI (%)	CS	CCI (%)
Bohol	113.7	I	44.8	I	98.5
		II	41.0	II	66.7
		III	39.3	III	51.2
		IV	39.3	IV	65.8
Cebu	145.8	I	31.4	I	66.7
		II	28.0	II	40.0
		III	26.5	III	28.4
		IV	26.5	IV	39.3
Negros Oriental	271.6	I	139.9	I	266.0
		II	128.9	II	188.5
		III	123.9	III	147.0
		IV	123.9	IV	186.3
Siquijor	264.5	I	80.4	I	182.2
		II	72.5	II	118.3
		III	69.1	III	87.8
		IV	69.1	IV	116.6

Crop Stage (CS) highlighted in black is the dominant stage during the month of September

Rice CS:
(I) Nursery
(II) Vegetative
(III) Reproductive
(IV) Ripening

Corn CS:
(I) Establishment
(II) Vegetative
(III) Reproductive
(IV) Maturity



CCI Category:
Inadequate
Sufficient
Excess

Regional Summary



Sufficient amount of rainfall was noted at the following: corn crops specifically those at the prevailing vegetative stage in the provinces of Bohol, Negros Oriental, and Siquijor; and rice crops at the expected harvesting and/ or pre-planting stage for the aforementioned provinces. On the other hand, inadequate amount of rainfall was noted in the province of Cebu for both rice and corn crops, and Bohol for rice crops at their prevailing stage.

Furthermore, the three-month accumulated rainfall (SPEI3) indicates generally near-normal to slightly wetter conditions on the central portion of Cebu and some portions in Bohol. While certain parts of Negros Oriental experienced slightly drier conditions. Apart from Negros Oriental, where the maximum rainfall recorded was 250 mm, the extreme rainfall indices (RX1day and RX5day) indicate no significant extreme rainfall that could have had a negative effect on crops in the area.


NDVI



SPEI3 (Jul-Aug-Sep)



RX1day



RX5day

